

IRAN STEPS Survey 2008

Fact Sheet 1387

This STEPS survey of chronic disease risk factors in Iran was carried out from 5/2008 to 6/2008. Iran carried out Step 1, Step 2; Socio demographic and behavioral information was collected in Step 1. Physical measurements such as height, weight and blood pressure were collected in Step 2.

The STEPS survey in I ran was a population-based survey of adults aged 15-64. A cluster sample design was used to produce representative data for that age range in I ran. A total of 30000 adults participated in the I ran STEPS survey. A repeat survey is planned for 2009.

Results for adults aged 15-64 years (incl. 95% CI) (adjust if necessary)	Both Sexes	Males	Females
Step 1 Tobacco Use			
Percentage who currently smoke tobacco daily	11.13 [10.58,11.71]	21.19 [20.66,21.74]	0.81 [0.71,0.93]
For those who smoke tobacco daily			
Average age started smoking (years)	20.67 [19.89,21.45]	19.45 [19.03,19.87]	21.91 [20.40,23.43]
Average years of smoking	13.15 [12.49,13.81]	14.48 [14.16,14.79]	11.79 [10.44,13.13]
Percentage smoking manufactured cigarettes			
For smokers of manufactured cigarettes			
Mean number of manufactured cigarettes smoked per day	9.14 [8.51,9.78]	12.29 [11.69, 12.90]	5.91 [4.80,7.03]
Step 1 Alcohol Consumption		1	
Percentage of abstainers (who did not drink alcohol in the last year)			
Percentage of current drinkers (who drank alcohol in the past 30 days)			
For those who drank alcohol in the last 30 days			
Percentage of women who had 4 or more drinks on any day in the last week			
Percentage of men who had 5 or more drinks on any day in the last week			
Percentage who drank alcohol on 4 or more days in the last week			
Step 1 Fruit and Vegetable Consumption (in a typical week)			
Mean number of servings of fruit consumed per day	1.44 [1.38,1.49]	1.41 [1.35,1.47]	1.46 [1.40,1.52]
Mean number of servings of vegetables consumed per day	1.40 [1.31,1.48]	1.34 [1.26,1.42]	1.46 [1.36,1.56]
Percentage who ate 5 or more combined servings of fruit & vegetables per day	11.75 [10.49,13.13]	11.15 [10.46,11.89]	12.35 [11.64,13.1]
Step 1 Physical Activity			
Percentage with low levels of activity (defined as <600 MET-minutes/week)	38.1 [36.8,39.42]	26.6 [25.86,27.36]	49.88 [49.02,50.75]
Median time spent in work-related physical activity per day (minutes)	0	4.28	0
Median time spent in transport-related physical activity per day (minutes)	15	17.14	11.42
Median time spent in recreational physical activity per day (minutes)	0	0	0



IRAN STEPS Survey 2008

Fact Sheet 1387

Results for adults aged 15-64 years (incl. 95% CI) (adjust if necessary)	Both Sexes	Males	Females
Step 2 Physical Measurements			•
Mean body mass index - BMI (kg/m²)	24.91	24.16	25.68
	[24.81,25.02]	[24.04,24.28]	[25.53,25.83]
Percentage who are overweight or obese (BMI ≥ 25 kg/m²)	44.25	38.78	49.85
	[43.26,45.25]	[38.17,39.4]	[49.14,50.57]
Percentage who are obese (BMI ≥ 30 kg/m²)	14.86	9.68	20.17
	[14.17,15.58]	[9.265,10.12]	[19.69,20.66]
Average waist circumference (cm)	84.33	85.18	83.47
	[83.99,84.68]	[84.78,85.58]	[83.01,83.93]
Mean systolic blood pressure - SBP (mmHg)	118.41	121.71	115.02
	[118.05,118.76]	[121.24,122.17]	[114.56,115.49]
Mean diastolic blood pressure - DBP (mmHg)	77.08	77.32	76.84
	[76.81, 77.35]	[76.96,77.67]	[76.52,77.17]
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg)	16.82	16.55	17.10
	[16.12,17.55]	[16.05,17.07]	[16.69,17.51]
Percentage with raised BP (SBP ≥ 170 and/or DBP ≥ 100 mmHg)	6.50	5.08	7.95
	[6.13,6.88]	[4.85,5.32]	[7.66,8.25]
Step 3 Biochemical Measurements			
Mean fasting blood glucose (mmol/L)			
Percentage with raised blood glucose (≥ 7.0 mmol/L)			
Mean total blood cholesterol (mmol/L)			
Percentage with raised total cholesterol (≥ 5.2 mmol/L)			
Percentage with raised total cholesterol (≥ 6.5 mmol/L)			
Summary of combined risk factors	<u> </u>		
 current daily smokers less than 5 servings of fruits & vegetables per day Low level of activity (<600 MET -minutes) overweight or obese (BMI ≥ 25 kg/m²) raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg) 			
Percentage with low risk (i.e. none of the risk factors included above)	3.19	3.58	2.78
	[2.69,3.77]	[3.27,3.92]	[2.50,3.10]
Percentage with raised risk (i.e. at least three of the risk factors included above) distributed in the age groups below			
Percentage with raised risk, aged 15 to 44 years old	22.47	20.98	24.00
	[21.47,23.49]	[20.37,21.6]	[23.3,24.72]
Percentage with raised risk, aged 45 to 64 years old	50.15	45.84	54.48
	[48.48,51.81]	[44.69,46.99]	[53.43,55.52]

For additional information, please contact:

Iran STEPs focal point; F.Asgari, <u>f-sgari@health.gov.ir</u>, <u>asgarifcdc@yahoo.com</u>
NCD Surveillance Risk Factor office: <u>ncdsurf@health.gov.ir</u>, <u>ncd_surveillance@yahoo.com</u>

For additional information, please contact: STEPS country focal point [name, email addresses]